

THE ASTHMA JOURNAL OF NEW ZEALAND

asthma update

May/June 2004

SPECIAL FEATURE:

BOILING A FROG – YOU & YOUR ASTHMA

COPD – DON'T LET THIS CONDITION
BE THE WINTER CINDERELLA

TREATMENT OF ASTHMA WITH
LIPID EXTRACT OF NEW ZEALAND
GREEN-LIPPED MUSSEL: A CLINICAL TRIAL

ASTHMA & SMOKING



\$4.50 (incl. GST)

This issue is proudly sponsored by:



Boehringer
Ingelheim



9 771368 254725 >



TREATMENT OF ASTHMA WITH LIPID EXTRACT OF NEW ZEALAND GREEN-LIPPED MUSSEL: A RANDOMISED CLINICAL TRIAL

A. EMEL'YANOV, G. FEDOSEEV, O. KRASNOSCHKOVA,
A. ABULIMITY, T. TRENDELEVA, P.J. BARNES

Eur Respir J 2002; 20: 1-5
DOI: 10.1183/09031936.00.00000000
Printed in UK – all rights reserved
Copyright ©ERS Journals Ltd 2002
European Respiratory Journal
ISSN 0903-1936

TREATMENT OF ASTHMA WITH LIPID EXTRACT OF NEW ZEALAND GREEN-LIPPED MUSSEL.



Treatment of asthma with lipid extract of New Zealand green-lipped mussel: A randomised clinical trial. A. Emelyanov, G. Fedoseev, O. Krasnoschekova, Medical University, St-Petersburg, A. Abulimity, T. Trendeleva, P.J. Barnes. ©ERS Journals Ltd 2002.

ABSTRACT: Asthma is a chronic inflammatory disease of the airways mediated at least in part by leukotrienes and other lipid mediators. Experimental studies have shown that lipid extract of New Zealand Green-lipped mussel, *Perna canaliculus*, is effective in inhibiting 5' – Lipoxygenase and cyclo-oxygenase pathways responsible for production of eicosanoids, including leukotrienes and prostaglandins. The aim of this study was to assess its effect on symptoms, peak expiratory flow (PEF) and hydrogen peroxide (H_2O_2) in expired breath condensate as a marker of airway inflammation in patients with steroid-naïve atopic asthma in a double-blind randomised, placebo-controlled clinical trial.

Forty six patients with atopic asthma received two capsules of lipid extract (Lyprinol®) or placebo *b.i.d.* for 8 weeks. Each capsule of lipid extract contained 50 mg ω -3 polyunsaturated fatty acids and 100 mg olive oil, whereas placebo capsules contained only 150 mg olive oil.

There was a significant decrease in daytime wheeze, the concentration of exhaled H_2O_2 and an increase in morning PEF in the lipid extract group compared to the placebo group. There were no significant side-effects.

The authors conclude that lipid extract of New Zealand green-lipped mussel may have some beneficial effect in patients with atopic asthma. *Eur Respir J* 2002; 20: 1-5.

Asthma is a chronic inflammatory disease of the airways mediated, at least in part, by leukotrienes and other lipid mediators. Experimental studies have shown that lipid

Table 1. – Characteristics of patients with atopic asthma treated with Lyprinol and placebo

CHARACTERISTICS OF PATIENTS	PLACEBO	LYPRINOL
Patients	23	23
Sex M-F	5:18	7:16
Age yrs	37.4±2.2	40.8±2.4
Duration of asthma yrs	4.5±0.9	7.32±1.2
FEV ₁ % pred	90.4±3.0	82.9±4.2
Concomitant medications	Short-acting inhaled β_2 -agonists as required	Short-acting inhaled β_2 -agonists as required

Data are expressed as mean±SEM. M: male; F: female. FEV₁: forced expiratory volume in one second. Differences between two groups of asthmatic patients are not significant.

mussel (*Perna canaliculus*) is effective at inhibiting 5' –lipoxygenase and cyclo-oxygenase pathways, which are responsible for the production of eicosanoids, including leukotrienes and prostaglandins (1-3). The lipid extract Lyprinol is rich in eicosapentaenoic acid and docosahexanoic acid, ω -3 fatty acids that inhibit the metabolism of arachidonic acid (4). This extract is effective in reducing pain, swelling and stiffness and in improving the functional index in patients with rheumatoid arthritis and osteoarthritis (5, 6). Its efficacy in the treatment of airway inflammation is unknown.

The aim of this study is to assess the effect of lipid extract of New Zealand green-lipped mussel on symptoms, peak expiratory flow (PEF) and hydrogen peroxide (H_2O_2) in expired breath condensate as a marker of airway inflammation in patients with atopic asthma in double-blind randomised, placebo-controlled clinical trial.

Methods

Patients

Volunteers aged between 18-56 yrs with mild-to-moderate atopic asthma were recruited in the out-patients department of the Hospital Therapeutic Clinic, Pavlov Medical University, St. Petersburg, Russia (table 1). Patients were diagnosed according to the American Thoracic Society definition of asthma (7). Diagnosis was based upon clinical history, reversibility of forced expiratory volume in one second (FEV₁) of > 15% and diurnal variability of PEF of > 20%. Atopic status was assessed by positive skin-prick test (>3 mm) to common inhalant allergens (house-dust mite, animal allergens, pollen). All patients were sensitised to house-dust mite. Severity of asthma was classified according to the National Institutes of Health/World Health

Organization (NIH/WHO) guidelines (8). Briefly, subjects with mild asthma had symptoms twice a week or less often with FEV₁ \geq 80% of predicted and used inhaled short-acting β_2 -agonists for symptom relief. Patients with moderate asthma had daily symptoms and an FEV₁ between 60 and 80% pred, and used inhaled short-acting β_2 -agonists daily. Exclusion criteria were use of inhaled steroids within the previous 4 weeks, oral steroid treatment within the previous 3 months, current or exsmokers, clinically significant heart, renal, liver and intestinal disorders, women of childbearing potential not using adequate contraception. Only patients fully co-operative in all procedures were considered for inclusion in the run-in period of the trial.

The study was approved by the Local Ethics Committee and all participants gave written consent.

Study design

The study was a single-centre, double-blind, parallel group, randomised, placebo-controlled trial. Patients who fulfilled the inclusion criteria at the initial visit entered a 2-week observation run-in period, in which regular treatment was stopped and short-acting β_2 -agonists (salbutamol or terbutaline) as required were the only medications allowed. Asthma symptoms and use of rescue medications were recorded in a diary. Patients were excluded if they received inhaled steroids or were hospitalised for asthma during the run-in period.

After the run-in period patients were randomly assigned to receive either lipid extract of New Zealand green-lipped mussel, two capsules twice daily or matching placebo for a period of 8 weeks. Each lipid extract capsule contained 50 mg ω -3 polyunsaturated fatty acids (eicosapentaenoic acid and docosahexaenoic

Continued on page 16

TREATMENT OF ASTHMA WITH LIPID EXTRACT OF NEW ZEALAND GREEN-LIPPED MUSSEL.

... from page 15

acid) and 100 mg olive oil (Lyprinol®, Mac Lab, Melbourne Australia). Each identical placebo capsule contained 150 mg olive oil only. Randomisation was computer-generated in balanced blocks of two treatment regimens. Inhaled salbutamol delivered via pressurised metered-dose inhaler or terbutaline delivered by dry powder inhaler (Turbuhaler) were used as rescue medications throughout the study. No other medications were allowed. All short-acting inhaled β_2 -agonists were stopped at least 6 h before baseline FEV₁ and exhaled H₂O₂ measurements.

Measurements

Subjects kept a twice-daily diary card throughout the study. They were provided with individual peak-flow meters (Vitalograph Ltd, Buckingham, UK). The best of three peak flow values were recorded before taking the morning and evening study medications. Patients recorded daytime wheeze, nocturnal awakenings from asthma, daily use of short acting inhaled β_2 -agonists and compliance with the study medications. Asthma symptoms were ranked on a scale from 0 (none) to 3 (very uncomfortable and interferes with normal activities). At the beginning and the end of the study, subjects attended the research clinic for measurement of FEV₁, which was measured by dry spirometry (Vitalograph Ltd). The concentration of exhaled H₂O₂ was measured at the beginning of the

study, and at 4 and 8 weeks of treatment. Study medications and inhaled short-acting β_2 -agonists were withheld for at least 6 h before spirometry and exhaled H₂O₂ measurements.

Expired breath condensate was collected using a glass condensing device, which was placed in a large chamber with ice as previously described (9). After mouth rinsing, subjects breathed tidally through a mouthpiece for 20 mins while wearing a nose-clip. The mouthpiece was also used as a saliva trap. The volume of condensate was 2-4 mL and H₂O₂ was assayed immediately. H₂O₂ was measured by using a colorimetric assay as described previously (10). Briefly, 100 μ L of condensate was mixed with 100 μ L of 3,3',5,5' tetramethylbenzidine in 0.42 M citrate buffer, pH 3.8, and 10 μ L of horseradish peroxidase (52.5 U.m⁻¹). The samples were incubated at room temperature for 20 min and the reaction stopped by the addition of 10 μ L 18 N sulphuric acid. The reaction product was measured spectrophotometrically (Model 46; Lomo Inc., St-Petersburg, Russia) at 450 nm. A standard curve of H₂O₂ was performed for each assay.

Safety of the treatment

At each visit subjects were specially asked for any adverse events and health problems that may have occurred. All unusual signs and symptoms were recorded for further

"The data from this study show that a stabilised lipid extract of New Zealand green-lipped mussel, *Perna canaliculus* (Lyprinol), reduces daytime wheeze and improves morning PEF in steroid-naïve patients with atopic asthma compared to placebo."

consideration. The safety of the treatment was evaluated by monitoring blood pressure, concentrations of creatinine, bilirubin, activity of liver transaminases and alkaline phosphatase in serum at clinical visits.

Analysis of data

Paired and unpaired two-tailed t-tests were used for statistical analysis. A p-value <0.05 was considered significant, and the data are expressed as mean \pm SEM.

Results

Subjects

Of 60 volunteers recruited to the study, 46 fulfilled the entry criteria and were randomised to treatment. The most common reason for withdrawal before randomisation was deterioration in asthma control (14 subjects). These patients had a lower FEV₁ (70.3 \pm 2.1% pred) than those who were randomised (87.60 \pm 2.5, p < 0.001). All randomised patients completed the treatment and were included in the statistical analysis. Thirty six of them had mild asthma and 10 patients had moderate asthma.

Compliance

The mean self-recorded compliance with the two-times daily study medications (Lyprinol or placebo) was 95% and 93%, respectively.

Asthma symptoms and bronchodilator use

Mean daytime wheeze was significantly reduced in patients treated with Lyprinol compared to those treated with placebo (table 2). There were no significant differences between the two treatment groups in respect to nocturnal awakening and use of short-acting β_2 -agonists (table 2).



Lung function tests

Mean FEV₁ and evening PEF during the treatment with Lyprinol did not significantly differ from those during the treatment with placebo. However, mean morning PEF was significantly higher during the treatment with Lyprinol than with placebo (table 2).

Exhaled hydrogen peroxide

Mean concentrations of exhaled H₂O₂ were significantly reduced during the treatment with Lyprinol compared to placebo (table 2).

Safety of the treatment

One patient treated with Lyprinol and one patient treated with placebo had skin itch. One patient in the Lyprinol group and two patients in the placebo group complained of a metallic taste. There were no significant changes in blood pressure, or in concentrations of serum creatinine, bilirubin, liver transaminases or alkaline phosphatase in either group of patients (data not shown).

Discussion

The data from this study show that a stabilised lipid extract of New Zealand green-lipped mussel, *Perna canaliculus* (Lyprinol), reduces daytime wheeze and improves morning PEF in steroid-naïve patients with atopic asthma compared to placebo.

There were no differences between patients treated with Lyprinol and placebo with respect to FEV₁. It should be noted that most of the patients had mild asthma and FEV₁>80% pred and thus there was little room for improvement in this measurement.

Lyprinol was well tolerated and had very few side effects. The findings of the present study are in agreement with a previously published study (1).

The mechanism by which lipid extract of New Zealand green-lipped mussels exerts its beneficial effect remains to be elucidated. Lyprinol inhibits leukotriene production by stimulated human polymorphonuclear leukocytes *in vitro* (2). Much of this activity was associated with its content of ω -3 polyunsaturated fatty acids (eicosapentaenoic acid and docosahexaenoic acid) and antioxidants such as carotenoids. ω -3 polyunsaturated acids inhibit arachidonate metabolism by 5'-lipoxygenase and cyclooxygenase (6), thus reducing the formation of leukotrienes and prostaglandins. Dietary

Table 2. – Efficacy results

Parameters	Placebo	Lyprinol	p-value
Patients n	23	23	
Daytime wheeze			
Baseline	1.41±0.42	2.27±0.36	0.131
4 week	1.23±0.26	1.07±0.23	0.637
Δ	-0.17±0.33	-1.20±0.24	0.023
8 week	1.29±0.17	0.73±0.21	0.041
Δ	-0.11±0.43	-1.53±0.42	0.026
Night awakenings			
Baselines	0.69±0.24	1.09±0.16	0.195
4 week	0.54±0.18	0.45±0.16	0.737
Δ	-0.15±0.27	-0.64±0.20	0.184
8 week	0.61±0.14	0.36±0.20	0.308
Δ	-0.08±0.29	-0.73±0.19	0.085
Usage of β₂-agonists puffs-day¹			
Baseline	2.17±0.64	3.17±0.56	0.252
4 week	1.92±0.43	1.92±0.78	1.0
Δ	-0.25±0.52	-1.25±0.55	0.202
8 week	2.00±0.48	1.58±0.42	0.253
Δ	-0.17±0.38	-1.59±0.60	0.022
FEV₁ % pred			
Baseline	92.32±2.9	82.89±4.2	0.073
8 week	90.53±3.2	82.87±3.6	0.124
Δ	-1.79±4.4	-0.02±2.85	0.708
Morning PEF L·min⁻¹			
Baseline	384.3±21.5	361.3±17.4	0.409
28 day	376.0±26.0	392.2±18.9	0.619
Δ	-8.3±9.4	30.9±13.6	0.022
56 day	350.9±21.3	408.3±18.7	0.049
Δ	-33.4±6.2	47.0±11.7	0.00001
Evening PEF L·min⁻¹			
Baseline	399.6±16.7	375.4±18.2	0.333
4 week	395.9±23.0	404.8±17.1	0.758
Δ	-3.7±11.4	29.4±11.5	0.047
8 weeks	403.9±18.3	406.5±19.7	0.923
Δ	4.3±10.3	31.1±14.6	0.136
Exhaled H₂O₂ μM			
Baseline	0.12±0.04	0.11±0.02	0.706
4 week	0.11±0.02	0.07±0.01	0.072
Δ	-0.01±0.03	-0.04±0.02	0.509
8 weeks	0.16±0.04	0.05±0.01	0.022
Δ	0.04±0.05	-0.06±0.01	0.0001

Data are expressed as mean±SEM. Δ: change from baseline value; FEV₁: forced expiratory volume in one second; PEF: peak expiratory flow; H₂O₂: hydrogen peroxide. Asthma symptoms were ranked on a scale from 0 (none) to 3 (very uncomfortable and interferes with normal activities).

TREATMENT OF ASTHMA WITH LIPID EXTRACT OF NEW ZEALAND GREEN-LIPPED MUSSEL.

... from page 17

supplementation with ω -3 polyunsaturated fatty acids can suppress synthesis of interleukin-1 and tumour necrosis factor- α by mononuclear cells (11). Epidemiological studies in Inuit, Dutch (12), American (13), Australian (14) and Japanese populations (15) have shown that a high dietary intake of ω -3 polyunsaturated fatty acids is associated with lower incidence of cardiovascular disease and inflammatory diseases, such as asthma and rheumatoid arthritis. Dietary supplementation with fish oil rich in the ω -3 polyunsaturated fatty acids over a 10-month period is beneficial in children with asthma following a strictly controlled diet and controlled allergen exposure (16).

Leukotrienes are mediators of airway inflammation in asthma. They induce bronchoconstriction and airway smooth muscle proliferation (17, 18), stimulate mucus secretion (19), increase microvascular permeability and cause infiltration of inflammatory cells such as eosinophils and neutrophils into the airways (20, 21). These cells release superoxide anions, which then undergo spontaneous or enzyme-catalysed dismutation to form H_2O_2 . H_2O_2 is a highly reactive oxygen species involved in cellular injury via further reactions leading to hydroxyl radical and lipid peroxidation products (22). Increased production of free radicals occurs in airway inflammation, and H_2O_2 is detectable in exhaled air (23). Several studies have shown that H_2O_2 in expired breath condensate may be a nondirect marker of oxidative stress and airway inflammation in asthmatic patients (24, 25). Exhaled H_2O_2 levels have previously been related to the eosinophil differential counts in induced sputum and activity of peripheral neutrophils in asthmatic patients (25, 26). Therefore, elevated concentrations of H_2O_2 , may result from an enhanced number and activity of inflammatory cells in the airways.

In the present study, the authors found that elevated concentrations of exhaled H_2O_2 were significantly decreased during treatment with Lyprinol compared to placebo. This may result from a reduction in the number of inflammatory cells in asthmatic airways because of decreased synthesis of leukotrienes and may also be due to the antioxidants such as β -carotene contained in Lyprinol.

In conclusion, this study shows that a lipid extract of New Zealand green-lipped mussel may have some beneficial effects in patients with steroid-naïve atopic asthma. Further clinical studies are needed to determine whether it may be useful as a supplement to other medications in the treatment of asthmatic patients.



REFERENCES

- Whitehouse MW, Macrides TA, Kalfatis NK, Betts WH, Haynes DR, Broadbent J. Anti-inflammatory activity of a lipid fraction (Lyprinol) from the New Zealand green-lipped mussel. *Inflammopharmacology* 1997; 5: 237-246.
- Rainsford KD, Whitehouse MW. Gastroprotective and anti-inflammatory properties of green-lipped mussel (*Perna canaliculus*) preparation. *Arzn Forsch* 1980; 30: 2128-2132.
- Halpern G-M. Anti-inflammatory effects of a stabilized lipid extract of *Perna Canaliculus* (Lyprinol®). *Allerg Immunol* 2000; 32: 272-278.
- Sinclair AJ, Murphy KJ, Li D. Marine lipids: overview "news insights and lipid composition of Lyprinol" *Allerg Immunol (Paris)* 2000; 32: 261-271.
- Gibson SLM, Gibson RG. The treatment of arthritis with lipid extract of *Perna canaliculus*: a randomised trial. *Compl Ther Med* 1998; 6: 122-126.
- Gibson RG, Gibson S, Conway V, Chappel D. *Perna canaliculus* in the treatment of arthritis *Practitioner* 1980; 224: 955-960.
- American Thoracic Society. Standards for the diagnosis and care of patients with chronic obstructive pulmonary disease (COPD) and asthma. *Am Rev Respir Dis* 1987; 136: 225-244.
- Diagnosis and classification. In: Global Initiative for asthma. NHLB/WHO Workshop Report. Publication number 95-3659. USA, National Heart Lung Blood Institute, 1995; pp. 48-61.
- Emelyanov A, Fedoseev G, Abuduani Abulimity, et al. Elevated concentrations of exhaled hydrogen peroxide in asthmatic patients. *Chest* 2002; 120: 1136-1139.
- Gallati H, Pracht I. Horseradish peroxidase: kinetic studies and optimisation of peroxide activity determination using the substrates (H_2O_2) and 3,3', 5,5'-tetramethylbenzidine. *J Clin Chem Clin Biochem* 1985; 23: 453-460.
- Endres S, Ghorbani R, Kelly VE, et al. The effect of dietary supplementation with ω -3 polyunsaturated fatty acids on the synthesis of interleukin-1 and tumor necrosis factor by mononuclear cells. *N Engl J Med* 1989; 320:265-271.
- Kromann N, Green A. Epidemiological studies in the Upernavik district, Greenland. Incidence of some chronic diseases 1950-1974. *Acta Med Scan* 1980; 208: 401-406.
- Schwartz J, Weiss ST. The relationship of dietary fish oil intake to level of pulmonary function in the first National Health and Nutritional Survey. *Eur Respir J* 1994; 7: 1821-1824.
- Peat JK, Salome CM, Woolcock AJ. Factors associated with bronchial hyperresponsiveness in Australian adults and children. *Eur Respir J* 1992; 5: 921-929.
- Satomi H, Monowa M, Hatano S, Iikura Y. An epidemiological study of the preventive effect of dietary fish on bronchial asthma. *Bull Inst Public Health* 1994; 43: 305-314.
- Nagakura T, Matsuda S, Shichijyo K, Sugimoto H, Hata K. Dietary supplementation with fish oil rich in ω -3 polyunsaturated fatty acids in children with bronchial asthma. *Eur Respir J* 2000; 16: 861-865.
- Piper PJ. Leukotrienes: potent mediators of airway constriction. *Int Arch Allergy Apopl Immunol* 1985; 76: Suppl. 1, 43-48.
- Wang CG, Du T, Xu LJ, Martin JG. Role of leukotriene D4 in allergen-induced increases in airway smooth muscle in rat. *Am Rev Respir Dis* 1994; 148:413-417.
- Marom Z, Shelhamer JH, Bach MK, Morton DR, Kaliner M. Slow-reacting substances leukotriene C4 and D4 increase the release of mucus from human airway in vitro. *Am Rev Respir Dis* 1982; 126: 449-451.
- Dahlen S-E, Bjorck, Heqvist P, et al. Leukotrienes promote plasma leakage and leukocyte adhesion in post-capillary venules: in vitro effects with relevance to the acute inflammatory response. *Proc Natl Acad Sci USA* 1981; 78: 3887-3891.
- Laitinen LA, Laitinen A, Haahtela T, Vilkkla V, Spur BW, Lee TH. Leukotriene E4 cause granulocyte infiltration into asthmatic airway. *Lancet* 1993; 341: 989-990.
- Mates JM, Sanchez-Jimenez F. Antioxidant enzymes and their implications in pathophysiologic processes. *Front Biosci* 1999; 4: 339-345.
- Kharitonov SA, Barnes PJ. Exhaled markers of pulmonary diseases. *Am J Respir Crit Care Med* 2001; 163: 1693-1722.
- Antzak A, Nowak D, Shariata B, Krol M, Plasecka G, Kurmanowska Z. Increased hydrogen peroxide and thiobarbituric acid-reactive products in expired breath condensate of asthmatic patients. *Eur Respir J* 1997; 9: 1235-1241.
- Horvath I, Donnelly LE, Kiss A, et al Combined use of exhaled hydrogen peroxide and nitric oxide in monitoring asthma. *Am J Respir Crit Care Med* 1998; 158: 1042-1046.
- Antzak A, Nowak D, Bialasiewicz P, Kasielski M. Hydrogen peroxide in expired air condensate correlates positively with early steps of peripheral neutrophil activation in asthmatic patients. *Arch Immunol Ther Exp (Warsz)* 1999; 47: 119-126.



"With Lyprinol[®], my breathing isn't a hurdle."

Jana Pittman, 400m hurdle world champion

"Before taking Lyprinol[®] I was often concerned when competing that my breathing could let me down. I can now go into competitions with confidence knowing that I can breathe easier."

Jana Pittman, 400m hurdle world champion

In many cases inflammation can affect airways and breathing passages. Clinical research has shown that a natural marine lipid extract contained in **Lyprinol[®]** can help reduce inflammation by blocking the major pathways in the body that produce inflammatory compounds.

So take Lyprinol[®]. And take a deep breath.

Blackmores is dedicated to helping Australians find more natural solutions to health issues and has been for over 60 years. Our supplements are made to high standards using quality vitamins, herbs and minerals.

Qualified health experts are on hand to answer questions on our award-winning website www.blackmores.com.au where you'll also find all the latest natural health information and news. Or call Blackmores **Advisory Service** on 1800 803 760.

Always read the label. Use only as directed.
If symptoms persist see your healthcare professional.

BLACKMORES[®]
THE BEST OF HEALTH

www.blackmores.com.au



For expert advice visit www.blackmores.com.au/lyprinol